

LA-UR-20-23978

Approved for public release; distribution is unlimited.

Title: A Brief Introduction to Gamma and Neutron Radiation

Author(s): Karpus, Peter Joseph

Intended for: Course material for DOD partners

Issued: 2020-06-01

Disclaimer:

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



A Brief Introduction to Gamma and Neutron Radiation


Pete Karpus

LA-UR-20-XXXXX

Objectives

- Describe gamma rays in plain English
- Give 3 reasons we measure gamma rays
- Describe shielding effect vs gamma energy
- List 4 components of the natural background
- Detail the main reason we care about neutrons
- List three good neutron moderators

Elements, Nuclides, and Isotopes



Constituents of the Atom: Protons, Neutrons, and Electrons

Elements: Defined by the number of protons in the nucleus

Isotopes: atoms from same element but with different number of neutrons

Nuclides (or Radionuclides): a more general term specifying element and atomic mass

^{12}C (Carbon-12):

- a) has 6 protons and 6 neutrons
- b) Atomic number (Z): # of protons
- c) Atomic Mass (A): # of protons + neutrons


Why Measure Gamma Rays?

- Electromagnetic interactions are well understood.
- Gamma rays can be highly penetrating.
- We can extract information passively.
- We can extract information particular to a **radionuclide**.
 - What nuclides are present?
 - How much is there? (limited by self shielding)

^{235}U or ^{238}U vs. just “U”

What are Gamma Rays?


- Gamma rays are electromagnetic (EM) radiation that come from the atomic nucleus
 - EM radiation = light (visible and otherwise)



Origin of Gamma rays


GAMMA-RAY SIGNATURE

Gammas from Alpha Decay



Gamma Ray Energies

- Units usually in kilo-electron volts (keV)
- $1 \text{ keV} = 1,000 \text{ eV}$
- $1 \text{ MeV} = 1,000,000 \text{ eV}$
- $1 \text{ TeV} = 1,000,000,000,000 \text{ eV}$
 - About what it takes to lift a mosquito 1 cm
- Roughly speaking:
 - Low energy: $\sim < 200 \text{ keV}$
 - Medium Energy: $\sim 200 - 1000 \text{ keV}$
 - High Energy: $> \sim 1000 \text{ keV}$



Gamma-Ray Interactions


- Gamma rays interact with matter in many ways.
- One of those ways is scattering.
- Scattering of gamma rays is analogous to the ricochet of a bullet.
- Scattering can make the data for the same source look very different in different measurement configurations.

Gamma-Ray Scattering

Gamma rays scatter everywhere:

- In the detector
- In the surroundings
- In the source itself

Every time gamma rays scatter they lose energy.




Scattering can complicate the data that you take in the field.

Gamma-Ray Attenuation (Shielding)

- Shielding reduces the intensity of full-energy gamma rays from a source
 - The lower the initial gamma energy the more effective the shield



The gammas from ^{137}Cs have a full energy of **662 keV**. They lose some of that energy as they scatter. Some are absorbed in the shield.




Shielding, Energy, & Intensity

- Gamma rays are (sort of) like bullets.

More high-energy bullets get through the shield than low energy bullets.




Although in this example the intensity is higher for the smaller caliber.




Why Measure Neutrons?

- Neutrons are a key signature of **plutonium**
- Neutrons can be highly penetrating.
- We can extract information passively.
- There are very few natural neutron sources compared to gamma sources




Common Origins of Terrestrial Neutron Radiation

Spontaneous Fission (SF)



Induced Fission



(α ,n) Reactions

alpha particle
(helium nuclei)
2 protons, 2 neutrons.



beryllium nucleus
4 protons, 5 neutrons.



carbon nucleus
6 neutrons, 6 protons



free neutron
1 neutron
(unstable)




Neutron Sources

- Reactors and fuel
- Particle Accelerators
- (α ,n) sources: used in research and industry
- Transuranics: spontaneous fission (Pu, Cf-252)
- Natural Neutron Background:
 - Generally from 'cosmic ray' spallation
 - Very weak signal compared with natural gamma-ray background

Almost all neutron sources are man-made!

Neutron Interactions

- Neutrons often can scatter and can be reflected
- Neutrons can also be captured by nuclei





Neutron Moderation


- Moderation is the slowing down (thermalization) of neutrons through collisions with *light* nuclei

Fast neutron



Materials with lots of hydrogen (H_2O , HDPE, HE, etc.) make good moderators.




Neutron Moderation



Neutrons on Heavy Nuclei:
Poor Moderation




Neutrons on Light Nuclei:
Good Moderation




Neutron Multiplication

- How many neutrons are produced for every “starter” neutron (e.g. from SF)?

Neutrons in the sphere have a better chance of inducing fission (producing more neutrons!) before they escape




Neutrons in the flat plane more likely escape before inducing fission




Reflection & Multiplication

Surrounding a neutron source with a reflecting material may send neutrons back into the material to induce more fissions.




Bare sphere




Gammas from Neutron Interactions

Neutron-Capture Gammas

Thermal neutron is absorbed by target nucleus.




Resulting nucleus emits a gamma ray.




Neutron-Scatter Gammas

Fast neutron scatters from target nucleus.






Resulting nucleus emits a gamma ray.



Natural Background Radiation

Gammas: From the Earth

Neutrons: From Space



U-238 Decay Series



Activity #1

List several things that can happen to neutrons in this picture



Activity #2

Which source is more penetrating “pound for pound”?
Which one is more *intense* (assuming the labeled activity is current) ?



122 keV
136 keV



1173 keV
1332 keV



Summary

- Gamma rays are a form of light emanating from atomic nuclei.
- Gamma rays are penetrating, give us information particular to a radionuclide, and can be measured passively
- Shielding preferentially attenuates lower-energy gamma rays
- The 4 main components of the natural background are:
 - gamma rays from K-40
 - gamma rays from uranium decay series
 - gamma rays from thorium decay series
 - neutrons from cosmic ray spallation
- Examples of good neutron moderators are water, HDPE, and HE as these all contain lots of hydrogen


Appendix

- The following concepts have been covered in the previously-delivered HP presentation but are included here for reference.

What is Radiation?


- When an atomic nucleus is unstable it emits something to get to its “ground state”.

Alpha Decay



This “something” is what we call “radiation”.

Radiation Types



Units of Radioactive Decay

- 1 disintegration per second = 1 Becquerel (Bq)
- 1 Curie (Ci) = 3.7×10^{10} Bq
- $1 \mu\text{Ci} = 1 \times 10^{-6}$ Ci (or 1 millionth of a Ci)
- Typical Check Sources:
 - ~ 1 to a few hundred μCi

